REPORT DOCUMENTATION PAGE

Form Approved OMB NO. 0704-0188

Public Reporting burden for this collection of information is estimated	d to average 1 hour per response, including the time for	r reviewing instructions, searching existing data sources, gathering and		
maintaining the data needed, and completing and reviewing the collect	ction of information. Send comment regarding this bur	den estimates or any other aspect of this collection of information,		
including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA				
22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188.) Washington, DC 20503.				

22202-4302, and to the Office of Management an			
1. AGENCY USE ONLY (Leave Blank)	2. REPORT DATE	3. REPORT TYPE	E AND DATES COVERED
	13 December 2000	Final Technical, 8	8/15/93 - 8/14/97
*			
4. TITLE AND SUBTITLE		5. FUNDING NUMB	
		DAAH04-93-G-03	396
Applications of New Concepts in	Scienticic Analysis to Atmospher	ic Studies	:
6. AUTHOR(S)			
James G. Brasseur			
7. PERFORMING ORGANIZATION N		8. PERFORMING O	
The Pennsylvania State Universi		gineering, REPORT NUMBE	ER .
123 Reber Building, University Pa	ark, PA 16802		
		· · · · · · · · · · · · · · · · · · ·	
9. SPONSORING / MONITORING AGI	ENCY NAME(S) AND ADDRESS(ES)	10. SPONSORING/	
]		AGENCY REPO	KI NUMBEK
U. S. Army Research Office	3		.
P.O. Box 12211		1 4203	2059.1-EV-AAS
Research Triangle Park, NO	27709-2211	/ TAYO 36	703 111-C1 MV2
Tresearen Triangie Turk, 100	32,703 2211	·	
11. SUPPLEMENTARY NOTES		L	
	:	one those of the author(s) and should	t not be construed as an official
		are those of the author(s) and should	I not be constitued as an official
Department of the Army position	i, policy or decision, unless so de	esignated by other documentation.	
12 a. DISTRIBUTION / AVAILABILIT	TY STATEMENT	12 b. DISTRIBUTION	ON CODE
L		1	
Approved for public release;	distribution unlimited.		**
			· .
Approved for public release; 13. ABSTRACT (Maximum 200 words)			· .
13. ABSTRACT (Maximum 200 words)		n constitute serious environment ha	zards
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar	pollutants in the atmosphere car	n constitute serious environment had	
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to	pollutants in the atmosphere car o the increased danger associate	d with extremely high instantaneous	s point
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed	pollutants in the atmosphere can the increased danger associate the local turbulence structure und	ed with extremely high instantaneous derlying statistical descriptions of tur	s point rbulent diffusion.
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the	pollutants in the atmosphere can the increased danger associate the local turbulence structure und entrainment of passive scalar ac	ed with extremely high instantaneous derlying statistical descriptions of tur coss a buoyancy-driven convective b	s point rbulent diffusion. ooundary layer
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using	pollutants in the atmosphere can the increased danger associate the local turbulence structure und entrainment of passive scalar act 1963 and 1283 large-eddy simula	ed with extremely high instantaneous derlying statistical descriptions of tul ross a buoyancy-driven convective b ation (LES). In the simulations, a un	s point rbulent diffusion. oundary layer iform concentration
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release.	pollutants in the atmosphere can to the increased danger associate the local turbulence structure und entrainment of passive scalar ac 1963 and 1283 large-eddy simulad in the stable layer above a CB	ed with extremely high instantaneous derlying statistical descriptions of ture ross a buoyancy-driven convective bation (LES). In the simulations, a un L with a strong capping inversion (9)	s point rbulent diffusion. noundary layer iform concentration 63 LES) and a CBL
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion.	pollutants in the atmosphere can the increased danger associate the local turbulence structure und entrainment of passive scalar act 1963 and 1283 large-eddy simulated in the stable layer above a CB 1 (1283 LES). The local turbulence	ed with extremely high instantaneous derlying statistical descriptions of turn ross a buoyancy-driven convective bation (LES). In the simulations, a un with a strong capping inversion (9) the structure in and around the entrain	s point rbulent diffusion. roundary layer iform concentration 63 LES) and a CBL nment layers were
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image.	pollutants in the atmosphere can the increased danger associated the local turbulence structure und entrainment of passive scalar act 1963 and 1283 large-eddy simulated in the stable layer above a CB 1 (1283 LES). The local turbulency ging, single-point statistical meas	ed with extremely high instantaneous derlying statistical descriptions of turn ross a buoyancy-driven convective bation (LES). In the simulations, a unusuith a strong capping inversion (9) the structure in and around the entrainstants and conditional statistics. Ame	s point rbulent diffusion. roundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra	pollutants in the atmosphere can the increased danger associated the local turbulence structure undentrainment of passive scalar act 1963 and 1283 large-eddy simulated in the stable layer above a CB 1 (1283 LES). The local turbulence ging, single-point statistical meastinment of top-down passive scal	ed with extremely high instantaneous derlying statistical descriptions of turn ross a buoyancy-driven convective bation (LES). In the simulations, a unblushed with a strong capping inversion (9) the structure in and around the entrainstance and conditional statistics. Amount to be more rapid with the weaker	s point rbulent diffusion. boundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment for	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undertrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulency sing, single-point statistical measurement of top-down passive scalar or above is affected by bounda	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unbut with a strong capping inversion (9) are structure in and around the entrainment of the more rapid with the weaker by layer growth, and that entrainments.	s point rbulent diffusion. boundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment for	pollutants in the atmosphere can the increased danger associated the local turbulence structure undentrainment of passive scalar act 1963 and 1283 large-eddy simulated in the stable layer above a CB 1 (1283 LES). The local turbulence ging, single-point statistical meastinment of top-down passive scal	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unbut with a strong capping inversion (9) are structure in and around the entrainment of the more rapid with the weaker by layer growth, and that entrainments.	s point rbulent diffusion. boundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment for	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undertrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulency sing, single-point statistical measurement of top-down passive scalar or above is affected by bounda	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unbut with a strong capping inversion (9) are structure in and around the entrainment of the more rapid with the weaker by layer growth, and that entrainments.	s point rbulent diffusion. boundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment for	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undertrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulency sing, single-point statistical measurement of top-down passive scalar or above is affected by bounda	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unbut with a strong capping inversion (9) we structure in and around the entrainment of the more rapid with the weaker by layer growth, and that entrainments.	s point rbulent diffusion. boundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment for	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undertrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulency sing, single-point statistical measurement of top-down passive scalar or above is affected by bounda	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unbut with a strong capping inversion (9) we structure in and around the entrainment of the more rapid with the weaker by layer growth, and that entrainments.	s point rbulent diffusion. boundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment for	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undertrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulency sing, single-point statistical measurement of top-down passive scalar or above is affected by bounda	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unbut with a strong capping inversion (9) we structure in and around the entrainment of the more rapid with the weaker by layer growth, and that entrainments.	s point rbulent diffusion. boundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment from in low vorticity downdraft region 14. SUBJECT TERMS	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undertrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulence ging, single-point statistical meast inment of top-down passive scalar on above is affected by boundations which transport scalar directions.	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unable with a strong capping inversion (9) he structure in and around the entrainsures and conditional statistics. Amount to be more rapid with the weaker ry layer growth, and that entrainmenty through the capping inversion.	s point rbulent diffusion. poundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion at was concentrated
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment from in low vorticity downdraft region 14. SUBJECT TERMS	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undertrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulency sing, single-point statistical measurement of top-down passive scalar or above is affected by bounda	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unable with a strong capping inversion (9) he structure in and around the entrainsures and conditional statistics. Amount to be more rapid with the weaker ry layer growth, and that entrainmenty through the capping inversion.	s point rbulent diffusion. poundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion at was concentrated
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment from in low vorticity downdraft region 14. SUBJECT TERMS	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undertrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulence ging, single-point statistical meast inment of top-down passive scalar on above is affected by boundations which transport scalar directions.	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unable with a strong capping inversion (9) he structure in and around the entrainsures and conditional statistics. Amount to be more rapid with the weaker ry layer growth, and that entrainmenty through the capping inversion.	s point rbulent diffusion. poundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion at was concentrated
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment from in low vorticity downdraft region 14. SUBJECT TERMS	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undertrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulence ging, single-point statistical meast inment of top-down passive scalar on above is affected by boundations which transport scalar directions.	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unable with a strong capping inversion (9) he structure in and around the entrainsures and conditional statistics. Amount to be more rapid with the weaker ry layer growth, and that entrainmenty through the capping inversion.	s point rbulent diffusion. coundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion at was concentrated
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment from in low vorticity downdraft region 14. SUBJECT TERMS	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undertrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulence ging, single-point statistical meast inment of top-down passive scalar on above is affected by boundations which transport scalar directions.	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unusual with a strong capping inversion (9) the structure in and around the entrainsures and conditional statistics. Amount of the more rapid with the weaker by layer growth, and that entrainment of the training of the training of the capping inversion.	s point rbulent diffusion. poundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion at was concentrated
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entral suggesting that entrainment from the in low vorticity downdraft region 14. SUBJECT TERMS dispersion, atmospheric bound 17. SECURITY CLASSIFICATION	pollutants in the atmosphere can of the increased danger associated the local turbulence structure undentrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulence ging, single-point statistical meast inment of top-down passive scalar on above is affected by boundarins which transport scalar directions which transport scalar directions.	ded with extremely high instantaneous derlying statistical descriptions of turioss a buoyancy-driven convective bation (LES). In the simulations, a unlaw with a strong capping inversion (9) we structure in and around the entrainsures and conditional statistics. Amount to be more rapid with the weaker by layer growth, and that entrainmenty through the capping inversion. The security classification	s point rbulent diffusion. coundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion at was concentrated
13. ABSTRACT (Maximum 200 words) The diffusion of passive scalar Recent studies have pointed to concentrations. We analyzed Specifically, we analyzed the (CBL) capping inversion using of passive scalar was release with a weak capping inversion analyzed using graphical image found that We found the entra suggesting that entrainment from low vorticity downdraft region 14. SUBJECT TERMS dispersion, atmospheric bounds.	pollutants in the atmosphere can the increased danger associated the local turbulence structure undentrainment of passive scalar acts 1963 and 1283 large-eddy simulated in the stable layer above a CB (1283 LES). The local turbulence ging, single-point statistical measinment of top-down passive scalar on above is affected by boundations which transport scalar directions which transport scalar directions.	ed with extremely high instantaneous derlying statistical descriptions of turnoss a buoyancy-driven convective bation (LES). In the simulations, a unusual with a strong capping inversion (9) the structure in and around the entrainsures and conditional statistics. Amount of the more rapid with the weaker by layer growth, and that entrainment of the training of the training of the capping inversion.	s point rbulent diffusion. poundary layer iform concentration 63 LES) and a CBL nment layers were ong other things we capping inversion at was concentrated

NSN 7540-01-280-5500

Standard Form 298 (Rev.2-89) Prescribed by ANSI Std. 239-18 298-102





MEMORANDUM OF TRANSMITTAL

U.S. Army Research Office ATTN: AMSRL-RO-BI (TR) P.O. Box 12211 Research Triangle Park, NC 27709-2211

☐ Reprint (Orig + 2 copies)	☐ Technical Report (Orig + 2 copies)	
Manuscript (1 copy)	Final Progress Report (Orig + 2 copies)	
	Related Materials, Abstracts, Theses (1 copy)	
CONTRACT/GRANT NUMBER: DAAH04-93-G-0396		
REPORT TITLE: Applications of New concepts in Scientific Analysis to Atmospheric Studies		
in formanded for your information		
is forwarded for your information.		
SUBMITTED FOR PUBLICATION TO (app	licable only if report is manuscript):	

Sincerely,

(James G Rocussium

Final Progress Report

ARO AASERT Grant DAAH04-93-G-0396 Applications of New Concepts in Scientific Analysis to Atmospheric Studies

Statement of problem studied

The diffusion of passive scalar pollutants in the atmosphere can constitute serious environment hazards. Recent studies have pointed to the increased danger associated with extremely high instantaneous point concentrations. We analyzed the local turbulence structure underlying statistical descriptions of turbulent diffusion. Specifically, we analyzed the entrainment of passive scalar across a buoyancy-driven convective boundary layer (CBL) capping inversion using 96³ and 128³ large-eddy simulation (LES). In the simulations, a uniform concentration of passive scalar was released in the stable layer above a CBL with a strong capping inversion (96³ LES) and a CBL with a weak capping inversion (128³ LES). The local turbulence structure in and around the entrainment layers were analyzed using graphical imaging, single-point statistical measures and conditional statistics.

Summary of most important results

We found the entrainment of top-down passive scalar to be more rapid with the weaker capping inversion suggesting that entrainment from above is affected by boundary layer growth. Locally, scalar penetration in the mixed layer was strongly influenced by the initial horizontal distribution of updrafts and downdrafts at the time of scalar release. The local dynamics of entrainment changed, however, after an initial period of high scalar flux lasting approximately one large-eddy turnover time. Entrainment then entered a quasi-steady asymptotic period with different entrainment behavior.

During the initial period, the local scalar flux in the immediate vicinity surrounding penetrative updrafts was negative and resulted form the turning motions of rising fluid downward. In the asymptotic state, by contrast, relatively little scalar entered the CBL in the immediate vicinity where the plumes were redirected downward by the capping inversion. Rather, most of the negative flux regions which contained positive scalar fluctuations coincided with downdrafts driven by the Rayleigh-Bénard cellular structures in the mixed layer. These far region downdrafts were the primary source of transport of scalar from above the capping inversion, they persisted independently of variations in the strength of neighboring updrafts, and they were the primary pathways for scalar entrainment. Localized mixing associated with large-horizontal-scale resolved vorticity to a large extend determined the local structure of positive scalar fluctuations found in these pathways, and this vorticity field appeared to determine the change in local structure of entrainment during the initial transition to the quasi-steady asymptotic entrainment state.

The description of a well-defined "entrainment layer" of smaller-scale eddies which transport scalar from above to below the mixed layer was not consistent with what was observed from the large-eddy simulations. Instead, entrainment was found to be concentrated in low vorticity downdraft regions which transport scalar directly through the capping inversion. Once the scalar was transported into the mixed layer, it was dispersed by local mixing processes. We find that top-down scalar diffusion is strongly influenced by the local structure of the mixed layer velocity and vorticity fields. The local velocity and vorticity structure appeared to determine the concentration levels and structure of the more intense positive scalar fluctuations within the mixed layer.

List of publications and technical reports

Cotter, John J. 1997 Scalar Entrainment through the Capping Inversion of the Atmospheric Boundary Layer. M.S. Thesis, Department of Mechanical Engineering, The Pennsylvania State University, University Park, PA.

Cotter, J.J., Brasseur, J.G., Khanna, S., Wyngaard, J.C. 1996 Scalar entrainment through the capping inversion of the atmospheric boundary layer. (abstract) *Bull. Amer. Phys. Soc.* 41(9): 1821

List of participating scientific personnel

John. C. Cotter III, M.S. student (supported on this grant)

Samir Khanna, Ph.D. student (supported on a related grant)

Brian Moquin, M.S. student (supported for 1 year on this grant)

John. C. Wyngaard, co-investigator with related grant

James G.Brasseur, pinciple investigator

Report on inventions

None.